



Restriction Map and Multiple Cloning Site (MCS) of pIRES2-EGFP Vector. Unique restriction sites are in bold. Note that the *Eco*47 III site has not been confirmed in the final construct.

Description:

pIRES2-EGFP contains the internal ribosome entry site (IRES; 1, 2) of the encephalomyocarditis virus (ECMV) between the MCS and the enhanced green fluorescent protein (EGFP) coding region. This permits both the gene of interest (cloned into the MCS) and the EGFP gene to be translated from a single bicistronic mRNA. pIRES2-EGFP is designed for the efficient selection (by flow cytometry or other methods) of transiently transfected mammalian cells expressing EGFP and the protein of interest. This vector can also be used to express EGFP alone or to obtain stably transfected cell lines without time-consuming drug and clonal selection.

EGFP is a red-shifted variant of wild-type GFP (3–5) which has been optimized for brighter fluorescence and higher expression in mammalian cells. (Excitation maximum = 488 nm; emission maximum = 507 nm.) EGFP encodes the GFPmut1 variant (6) which contains the double-amino-acid substitution of Phe-64 to Leu and Ser-65 to Thr. The coding sequence of the EGFP gene contains more than 190 silent base changes which correspond to human codon-usage preferences (7). Sequences flanking EGFP have been converted to a Kozak consensus translation initiation site (8) to further increase the translation efficiency in eukaryotic cells. The MCS in pIRES2-EGFP is between the immediate early promoter of cytomegalovirus ($P_{CMV\ IE}$) and the IRES sequence. SV40 polyadenylation signals downstream of the EGFP gene direct proper processing of the 3' end of the bicistronic mRNA. The vector backbone also contains an SV40 origin for replication in mammalian cells expressing the SV40 T antigen. A neomycin-resistance cassette (Neo'), consisting of the SV40 early promoter, the neomycin/kanamycin resistance gene of Tn5, and polyadenylation signals from the herpes simplex virus thymidine kinase (HSV TK) gene, allows stably transfected eukaryotic cells to be selected using G418. A bacterial promoter upstream of this cassette expresses kanamycin resistance in *E. coli*. The pIRES2-EGFP backbone also provides a pUC origin of replication for propagation in *E. coli* and an f1 origin for single-stranded DNA production. pIRES2-EGFP replaces (but is not derived from) the pIRES-EGFP Vector previously sold by BD Biosciences Clontech. pIRES2-EGFP is functionally similar to pIRES-EGFP; however, pIRES2-EGFP gives brighter EGFP fluorescence than the older vector. Note that the *Xba* I site at position

(PR29951; published 03 October 2002)



Clontech

United States/Canada
800.662.2566

Asia Pacific
+1.650.919.7300

Europe
+33.(0)1.3904.6880

Japan
+81.(0)77.543.6116

Clontech Laboratories, Inc.
A Takara Bio Company
1290 Terra Bella Ave.
Mountain View, CA 94043
Technical Support (US)
E-mail: tech@clontech.com
www.clontech.com

1987 is methylated in the DNA provided by BD Biosciences Clontech. If you wish to digest the vector with this enzyme, you will need to transform the vector into a *dam*⁻ host and make fresh DNA.

Use:

Genes inserted into the MCS should include the initiating ATG codon. pIRES2-EGFP and its derivatives can be introduced into mammalian cells using any standard transfection method. If required, stable transformants can be selected using G418 (9).

Location of features:

- Human cytomegalovirus (CMV) immediate early promoter: 1–589
Enhancer region: 59–465; TATA box: 554–560; Transcription start point: 583
C→G mutation to remove *Sac*I site: 569
- MCS: 591–665
- IRES sequence: 666–1250
- Enhanced green fluorescent protein (EGFP) gene
Kozak consensus translation initiation site: 1247–1257
Start codon (ATG): 1254–1256; Stop codon: 1971–1973
Insertion of Val at position 2: 1257–1259
GFPmut1 chromophore mutations (Phe-64 to Leu; Ser-65 to Thr): 1446–1451
His-231 to Leu mutation (A→T): 1948
- SV40 early mRNA polyadenylation signal
Polyadenylation signals: 2127–2132 & 2156–2161; mRNA 3' ends: 2165 & 2177
- f1 single-strand DNA origin: 2224–2679 (Packages the noncoding strand of EGFP.)
- Bacterial promoter for expression of Kan^r gene:
–35 region: 2741–2746; –10 region: 2764–2769
Transcription start point: 2776
- SV40 origin of replication: 3020–3155
- SV40 early promoter/enhancer
72-bp tandem repeats: 2853–2996; 21-bp repeats (3): 3000–3063
Early promoter element: 3076–3082
- Kanamycin/neomycin resistance gene: 3204–3998
G→A mutation to remove *Pst*I site: 3386; C→A (Arg to Ser) mutation to remove *Bss*H II site: 3732
- Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals: 4234–4252
- pUC plasmid replication origin: 4583–5226

Propagation in *E. coli*

- Suitable host strains: DH5 α , HB101, and other general purpose strains. Single-stranded DNA production requires a host containing an F plasmid such as JM101 or XL1-Blue.
- Selectable marker: plasmid confers resistance to kanamycin (30 μ g/ml) to *E. coli* hosts.
- *E. coli* replication origin: pUC
- Copy number: ~500
- Plasmid incompatibility group: pMB1/ColE1

References:

1. Jackson, R. J., *et al.* (1990) *Trends Biochem. Sci.* **15**:477–483.
2. Jang, S. K., *et al.* (1990) *J. Virol.* **62**:2636–2643.
3. Cormack, B., *et al.* (1996) *Gene* **173**:33–38.
4. Yang, T. T., *et al.* (1996) *Nucleic Acids Res.* **24**:4592–4593.
5. Haas, J., *et al.* (1996) *Curr. Biol.* **6**:315–324.
6. Jackson, R. J., *et al.* (1990) *Trends Biochem.* **15**:477–483.
7. Jang, S. K., *et al.* (1988) *J. Virol.* **62**:2636–2643.
8. Huang, M. T. F. & Gorman, C. M. (1990) *Nucleic Acids Res.* **18**(4):937–947.
9. Gorman, C. (1985). In *DNA cloning: A practical approach*, vol. II. Ed. D.M. Glover. (IRL Press, Oxford, U.K.) pp. 143–190.

Note: The attached sequence file has been compiled from information in the sequence databases, published literature, and other sources, together with partial sequences obtained by BD Biosciences Clontech. This vector has not been completely sequenced.

Notice to Purchaser

Use of BD Biosciences Clontech's Living Colors™ products containing DNA sequences coding for mutant *Aequorea victoria* green fluorescent protein (GFP) variants or proteins thereof requires a license from Amersham Biosciences under U.S. Patent Nos. 5,625,048; 5,777,079; 6,054,321 and other pending U.S. and foreign patent applications. In addition, certain BD Biosciences Clontech products are made under U.S. Patent No. 5,804,387 licensed from Stanford University.

Not-For-Profit research institutes or entities are granted an automatic license with the purchase of this product for use in non-commercial internal research purposes, the terms of which are disclosed in detail in the license that accompanies the shipment of this product. Such license specifically excludes the right to sell or otherwise transfer this product or its components to third parties.

For-Profit research institutes or entities must obtain a license from Amersham Biosciences. E-mail: gfp@amershambiosciences.com

Please contact BD Biosciences Clontech directly for any other assistance, including purchasing and technical support. All companies and institutions purchasing Living Colors™ products will be included in a quarterly report to Aurora Biosciences, as required by the BD Biosciences Clontech/Aurora Biosciences license agreement.

Use of the IRES sequence is covered by U.S. Patent #4,937,190 and is limited to use solely for research purposes. Any other use of the IRES sequence requires a license from Wisconsin Alumni Research Foundation.

This product is intended to be used for research purposes only. It is not to be used for drug or diagnostic purposes nor is it intended for human use. BD Biosciences Clontech products may not be resold, modified for resale, or used to manufacture commercial products without written approval of BD Biosciences Clontech.

© 2002, Becton, Dickinson and Company